arXiv:1406.6264v1 [math.GT] 24 Jun 2014

DEHN SURGERY AND SEIFERT SURFACE SYSTEM

MAKOTO OZAWA AND KOYA SHIMOKAWA

Abstract. For a compact connected 3-submanifold with connected boundary in the 3-sphere, we relate the existence of a Seifert surface system for a surface with a Dehn surgery along a null-homologous link. As its corollary, we obtain a refinement of the Fox's re-embedding theorem.

1. Introduction

Definition 1.1. Let M be a compact connected 3-manifold with connected boundary of genus g. A *spanning surface system* $\{F_i\}$ for M is a set satisfying the following.

(1) $\{F_i\}$ is a set of disjoint orientable surfaces properly embedded in M.
(2) $\{\partial F_i\}$ is a set of g disjoint loops $C_1, \dots, C_g$ which do not separate ∂M.

A spanning surface system $\{F_i\}$ for M is *completely disjoint* if $\{F_i\}$ is a set of g disjoint orientable surfaces.

Remark 1.2. We remark by [10, Corollary 1.4] that if M is a handlebody and $\{F_i\}$ is a completely disjoint spanning surface system for M, then there exists a meridian disk system $\{D_i\}$ for M such that $\partial F_i = \partial D_i$ for $i = 1, \dots, g$.

By a homological argument, we have the following proposition.

Proposition 1.3. *Any compact connected 3-submanifold with connected boundary in S^3 admits a spanning surface system.*

However, a compact connected 3-submanifold with connected boundary in S^3 does not always admit a completely disjoint spanning surface system.

Remark 1.4. We remark by [9] that there exists a compact connected 3-submanifold with connected boundary of genus 2 in S^3 which does not admit a completely disjoint spanning surface system.

Definition 1.5. Let S be a genus $g > 0$ closed surface in S^3, and put $S^3 = V \cup_S V'$. A *Seifert surface system* $(\{F_i\}, \{F'_i\})$ for S is a pair of sets satisfying the following.

(1) $\{F_i\}$ (resp. $\{F'_i\}$) is a spanning surface system for V (resp. V').
(2) $|C_i \cap C'_j| = \delta_{ij}$ for $i, j = 1, \dots, g$, where $\{\partial F_i\} = \{C_1, \dots, C_g\}$ and $\{\partial F'_i\} = \{C'_1, \dots, C'_g\}$

2010 *Mathematics Subject Classification.* 57M25.

Key words and phrases. link, Dehn surgery, Seifert surface.

The first author is partially supported by Grant-in-Aid for Scientific Research (C) (No. 23540105 and No. 26400097), The Ministry of Education, Culture, Sports, Science and Technology, Japan.

The second author is partially supported by Grant-in-Aid for Scientific Research (C) (No. 25400080), The Ministry of Education, Culture, Sports, Science and Technology, Japan.

A Seifert surface system $(\{F_i\},\ \{F_i'\})$ for S is *completely disjoint* if $\{F_i\}$ and $\{F_i'\}$ are completely disjoint.

Definition 1.6. Let $L = L_1 \cup \cdots \cup L_n$ be a link in S^3. Following [11], we say that L is a *reflexive link* if the 3-sphere can be obtained by a non-trivial Dehn surgery along L. In particular, if the surgery slope γ_i for L_i is $1/n_i$ for some integer n_i $(i = 1, \ldots, n)$, then we call the Dehn surgery a *$1/\mathbb{Z}$-Dehn surgery*.

Suppose that L is contained in a compact 3-submanifold M in S^3. We say that L is *null-homologous* in M if $[L] = 0$ in $H_1(M; \mathbb{Z})$, and that L is *completely null-homologous* in M if $[L_i] = 0$ in $H_1(M; \mathbb{Z})$ for $i = 1, \ldots, n$.

Theorem 1.7. *Let M be a compact connected 3-submanifold with connected boundary in S^3. Then the followings hold.*

1. *There exists a null-homologous link L in M, which is reflexive in S^3, such that a handlebody can be obtained from M by a $1/\mathbb{Z}$-Dehn surgery along L.*
2. *M admits a completely disjoint spanning surface system if and only if there exists a completely null-homologous link L in M, which is reflexive in S^3, such that a handlebody can be obtained from M by a $1/\mathbb{Z}$-Dehn surgery along L.*

We remark that in Theorem 1.7 (2), we can take a completely null-homologous reflexive link L so that it is disjoint from the completely disjoint spanning surface system.

Remark 1.8. Let M be a compact connected 3-submanifold of S^3 with connected boundary of genus g. Let $f : M \to X$ be a map onto a genus g handlebody X. We say that f is a *boundary preserving map* of M onto X if f is continuous and $f|_{\partial M}$ is a homeomorphism onto ∂X. We say that M is *retractable* if M can be retracted onto a wedge of g simple closed curves. If such a wedge can be chosen to be in ∂M, then M is called *boundary retractable*. Set $G = \pi_1(M)$ and define $G_1 = [G, G]$, $G_{n+1} = [G_n, G]$, $G_\omega = \bigcap_n G_n$. Then the following conditions are equivalent.

1. M admits a completely disjoint spanning surface system.
2. There exists a boundary preserving map from M onto a handlebody.
3. M is boundary retractable.
4. The natural map $\pi_1(\partial M) \to G/G_\omega$ is an epimorphism.

The equivalence between (1) and (2) was shown in [9, Theorem 2]. The equivalence between (2) and (3) was shown in [7, Theorem 3]. The equivalence between (3) and (4) was shown in [6, Theorem 2, 3].

Let M be a compact connected 3-submanifold of S^3. By Proposition 1.3, each component of the exterior of M admits a spanning surface system. If we adapt Theorem 1.7 (1) to the exterior of M, then we obtain a refinement of the Fox's re-embedding theorem as the following corollary.

Corollary 1.9 ([3, 16, 13])**.** *Every compact connected 3-submanifold M of S^3 can be re-embedded in S^3 so that the exterior of the image of M is a union of handlebodies.*

Remark 1.10. In relation with Remark 1.8, there is an another equivalence condition. Let M be a compact connected 3-submanifold of S^3 with connected boundary of genus 2. By Corollary 1.9, there exists a re-embedding of M so that its exterior is a genus 2 handlebody V. A handcuff graph shaped spine Γ of V is a *boundary spine* if its constituent link L_Γ is a boundary link that admits a pair of disjoint Seifert

surfaces whose interiors are contained in $S^3 - \Gamma$. A handlebody V is $(3)_S$-*knotted* if it does not admit any boundary spine. Then it was shown in [1, Theorem 3.10] that M admits a completely disjoint spanning surface system if and only if H is not $(3)_S$-knotted.

Theorem 1.7 also deduces the following corollary. It follows from (2) of the next corollary that every closed surfaces with completely disjoint Seifert surface systems can be related by $1/\mathbb{Z}$-Dehn surgeries along completely disjoint null-homologous reflexive links.

Corollary 1.11. *Let S be a closed surface in S^3 which separates S^3 into 3-submanifolds M and M' Then the followings hold.*

1. *There exist null-homologous links L and L' in M and M', which are reflexive in S^3, such that handlebodies can be obtained from M and M' by $1/\mathbb{Z}$-Dehn surgeries along L and L'.*
2. *S admits a completely disjoint Seifert surface system if and only if there exist completely null-homologous links L and L' in M, which are reflexive in S^3, and M' such that handlebodies can be obtained from M and M' by $1/\mathbb{Z}$-Dehn surgeries along L and L'.*

By Corollary 1.11 (1), we can obtain a Seifert surface system from a meridian-longitude disk system for the handlebodies by tubing along the null-homologous links.

Corollary 1.12. *Any closed surface in S^3 admits a Seifert surface system.*

Let M be a 3-manifold. Let $L \subset M$ be a submanifold with or without boundary. When L is 1 or 2-dimensional, we write $E(L) = M - int\, N(L)$. When L is of 3-dimension, we write $E(L) = M - int\, N(L)$.

2. Proof

Let V be a genus g handlebody in S^3, and $\{D_i\}$ be a meridian disk system for V. Since $V - \bigcup_i intN(D_i)$ is a 3-ball, there exists a spine Γ of V such that:

1. Γ consists of g loops $l_1, \ldots, l_g$ and g arcs $\gamma_1, \ldots, \gamma_g$ connecting l_i to a point x.
2. The point x is the center of the 3-ball $V - \bigcup_i intN(D_i)$, and which is homeomorphic to $N(x \cup \gamma_1 \cup \cdots \cup \gamma_g)$.
3. Each loop l_i is dual to D_i.

We call this spine Γ a *g-handcuff graph shaped spine* for V with respect to $\{D_i\}$.

Next, let $\{F_i\}$ be a set of orientable surfaces with boundary and without closed component. We say that $\{F_i\}$ is a *Seifert surface system* for Γ if $(\bigcup_i F_i) \cap \Gamma = \bigcup_i \partial F_i = \bigcup_i l_i$.

Lemma 2.1. *Any g-handcuff graph shaped spine in S^3 admits a Seifert surface system.*

Proof. We take a regular diagram of Γ such that $x \cup \gamma_1 \cup \cdots \cup \gamma_g$ has no crossing. Then we apply the Seifert's algorithm ([17]) to loops $l_1 \cup \cdots \cup l_g$ with arbitrary orientations, and obtain a Seifert surfaces $\{F_i'\}$ for the loops. □

The following lemma states that from any meridian disk system for a handlebody, we can obtain a Seifert surface system for the boundary of the bandlebody.



FIGURE 1. A g-handcuff graph shaped spine for V with respect to $\{D_i\}$

Lemma 2.2. *Let V be a genus g handlebody in S^3 with a meridian disk system $\{D_i\}$. Then there exists a spanning surface system $\{F_i\}$ for $E(V)$ such that $(\{D_i\},\{F_i\})$ is a Seifert surface system for ∂V.*

Proof. Let Γ be a g-handcuff graph shaped spine Γ for V with respect to $\{D_i\}$. By Lemma 2.1, Γ admits a Seifert surface system $\{F_i'\}$. The restriction $\{F_i'\}$ to $E(V)$ gives a spanning surface system, say $\{F_i\}$, for $E(V)$ such that $(\{D_i\},\{F_i\})$ is a Seifert surface system for ∂V. □

Let Γ be a g-handcuff graph shaped spine with a Seifert surface system $\{F_i\}$. We call the operation of (1) in Figure 2 a *band-crossing change* of $\{F_i\}$, and the operation of (2) in Figure 2 a *full-twist* of $\{F_i\}$. We remark that these operations can be obtained by a $1/\mathbb{Z}$-Dehn surgery along trivial links in the complement of $\{F_i\}$.



(1) a band-crossing change (2) a full-twist

FIGURE 2. A band-crossing change and a full-twist of $\{F_i\}$

Lemma 2.3. *Any g-handcuff graph shaped spine Γ with a Seifert surface system $\{F_i\}$ can be unknotted by band-crossing changes and full-twists of $\{F_i\}$.*

Proof. It is observed that Γ with $\{F_i\}$ can be transformed to be a "standard planar form" (cf. [8]) by the following operations.

(1) a band-crossing change of $\{F_i\}$
(2) a full-twists of $\{F_i\}$
(3) a crossing change between $\{F_i\}$ and $\{\gamma_i\}$



FIGURE 3. A g-handcuff graph shaped spine Γ with a Seifert surface system $\{F_i\}$, which is a "standard planar form"

(4) a crossing change among $\{\gamma_i\}$

However, the operations (3) and (4) can be exchanged for the operation (1). If Γ with $\{F_i\}$ has a standard planar form, then Γ is unknotted and this completes the proof. We remark that in a standard planar form, $l_1 \cup \cdots \cup l_g$ is the trivial link. □

Lemma 2.4. *Let L be a reflexive link in S^3 which is contained in a compact 3-submanifold M in S^3. Suppose that L in null-homologous (resp. completely null-homologous) in M. Then the core link L^* in the 3-submanifold M' obtained by a $1/\mathbb{Z}$-Dehn surgery along L is also null-homologous (resp. completely null-homologous) in M'.*

Proof. Suppose that L in null-homologous (resp. completely null-homologous) in M. Then L bounds a Seifert surface F (resp. completely disjoint Seifert surface) in M. Put $F^* = F \cap E(L)$. By a $1/\mathbb{Z}$-Dehn surgery, the meridian of the core link L^* intersects each component of ∂F^* in one point. This shows that F^* can be extended to a a Seifert surface (resp. completely disjoint Seifert surface) for L^* in M'. Thus L^* is also null-homologous (resp. completely null-homologous) in M'. □

Lemma 2.5. *Let V be a handlebody in S^3. Then the followings hold.*

(1) *∂V admits a Seifert surface system if and only if there exists a null-homologous link L in $E(V)$, which is reflexive in S^3, such that a handlebody can be obtained from $E(V)$ by a $1/\mathbb{Z}$-Dehn surgery along L.*
(2) *∂V admits a completely disjoint Seifert surface system if and only if there exists a completely null-homologous link L in $E(V)$, which is reflexive in S^3, such that a handlebody can be obtained from $E(V)$ by a $1/\mathbb{Z}$-Dehn surgery along L.*

Proof. (1) Suppose that there exists a null-homologous reflexive link L in $E(V)$ such that a handlebody, say W, can be obtained from $E(V)$ by a $1/\mathbb{Z}$-Dehn surgery along L. There exists a meridian disk system $\{D_i\}$ for W. Since L is null-homologous in $E(V)$, by Lemma 2.4, the core link L^* is also null-homologous in W. Therefore, we can obtain a Seifert surface system for $E(V)$ by tubing $\{D_i\}$ along L^*.

Conversely, suppose that ∂V admits a Seifert surface system $(\{F_i\}, \{F_i'\})$, where $\{F_i\}$ and $\{F_i'\}$ are spanning surface systems for V and $E(V)$. By Remark 1.2, we may assume that each F_i is a disk. Then there exists a g-handcuff graph shaped spine Γ and F_i' can be extended to a Seifert surface system for Γ. By Lemma 2.3, Γ with $\{F_i'\}$ can be unknotted, hence $E(V)$ is a handlebody, by band-crossing changes and full-twists of $\{F_i'\}$. This operations can be obtained by a $1/\mathbb{Z}$-Dehn surgery along a trivial link L in $\Gamma \cup \bigcup_i F_i'$. Since L is contained in $E(V) - \bigcup_i F_i'$, L is a null-homologous link in $E(V)$. Hence we obtain a null-homologous reflexive link L in $E(V)$ such that a handlebody can be obtained from $E(V)$ by a $1/\mathbb{Z}$-Dehn surgery along L.

(2) This can be proved by the argument similar to (1). □

Lemma 2.6. *Let S be a Heegaard surface in S^3 which decomposes S^3 into two handlebodies V and V'. Let $\{D_i\}$ be a meridian disk system for V. Then there exist a null-homologous reflexive link L' in V' which yields a handlebody V'' by a $1/\mathbb{Z}$-Dehn surgery on L' and a meridian disk system $\{D_i''\}$ for V'' such that $(\{D_i\}, \{D_i''\})$ is a completely disjoint Seifert surface system for S in $V \cup V''$.*

Proof of Lemma 2.6. We take a g-handcuff graph shaped spine Γ of V with respect to D_i. Since Γ can be unknotted by crossing changes, there exists a null-homologous reflexive link L' in V' such that after a $1/\mathbb{Z}$-Dehn surgery along L', every loops of Γ bound mutually disjoint disks. Therefore, a handlebody V'' obtained from V' by a $1/\mathbb{Z}$-Dehn surgery along L' admits a meridian disk system $\{D_i''\}$ so that $(\{D_i\}, \{D_i''\})$ is a completely disjoint Seifert surface system for S. □

Proof of Theorem 1.7. (1) We prove by an induction on the genus $g(\partial M)$. Since the 3-sphere does not contain an incompressible closed surface, there exists a compressing disk D for ∂M in S^3. We divide the proof into two cases.

Case 1: $D \subset M$
Case 2: $D \subset E(M)$

In Case 1, put $M' = M - int\, N(D)$. By the assumption of the induction, there exists a null-homologous reflexive link L' in M' such that handlebodies can be obtained from M' by a $1/\mathbb{Z}$-Dehn surgery along L'. This proves the theorem since M is obtained by adding 1-handle $N(D)$ to M'.

In Case 2, we take a maximal compression body W for ∂M in $E(M)$ [2]. If W is a handlebody (i.e. $W = E(M)$), then the theorem follows Lemma 2.2 and Lemma 2.5 (1). Otherwise as $g(\partial W) < g(\partial M)$, by the assumption of the induction, there exists a null-homologous reflexive link L' in each component of $E(M) - int\, W$ such that handlebodies can be obtained from the component by a $1/\mathbb{Z}$-Dehn surgery along L'. After these $1/\mathbb{Z}$-Dehn surgery, $E(M)$ is a handlebody. Therefore, again by Lemma 2.2 and Lemma 2.5 (1), there exists a null-homologous reflexive link L in M such that a handlebody can be obtained from M by a $1/\mathbb{Z}$-Dehn surgery along L. Finally, we recover the previous $1/\mathbb{Z}$-Dehn surgery on each component of $E(M) - int\, W$ to obtain the original $E(M)$.

(2) Suppose that there exists a completely null-homologous reflexive link L in M such that a handlebody can be obtained from M by a $1/\mathbb{Z}$-Dehn surgery along L. There exists a meridian disk system $\{D_i\}$ for the resultant handlebody. Since L is completely null-homologous in M, we can obtain a completely disjoint spanning surface system for M by tubing $\{D_i\}$ along L.

Conversely, suppose that M admits a completely disjoint spanning surface system $\{F_i\}$. In the following 3 steps, we convert M and $E(M)$ into two handlebodies V and V'' so that (V, V'') admits a meridian disk system $(\{D_i\}, \{D_i''\})$ with $\partial F_i = \partial D_i$.

Step 1: By (1) of this theorem, there exists a null-homologous link reflexive L in $E(M)$ such that a handlebody can be obtained from $E(M)$ by a $1/\mathbb{Z}$-Dehn surgery along L. Let V' the resultant handlebody obtained from $E(M)$ and note that $M \cup V'$ is again the 3-sphere.

Step 2: We note that there exists a degree one map from M to a handlebody V which sends each F_i to a meridian disk D_i of V and preserves the boundary of M (cf. [9, Theorem 2], [4, Theorem 5]). We naturally extend this degree one map to a degree one map $\phi : S^3 = M \cup V' \to X = V \cup V'$ as follows.

(1) V' is contained in X by an inclusion.
(2) Each F_i is sent to a meridian disk D_i of the handlebody $\phi(M) = V$.
(3) The remnant $M - \bigcup int\, N(F_i)$ is sent to the 3-ball $V - \bigcup int\, N(D_i)$.

Since $\phi_* : \pi_1(S^3) \to \pi_1(X)$ is surjective [5, Lemma 15.12], X is homeomorphic to S^3 [14, 15, 12].

Step 3: By Lemma 2.6, there exists a null-homologous reflexive link L' in V' and a meridian disk system $\{D_i''\}$ for a handlebody V'' obtained from V' by a $1/\mathbb{Z}$-Dehn surgery along L' such that $(\{D_i\}, \{D_i''\})$ is a completely disjoint Seifert surface system for (V, V'').

Since the degree one map ϕ is a boundary preserving map by the condition (1), $(\{F_i\}, \{D_i''\})$ is a completely disjoint Seifert surface system for (M, V''). By Lemma 2.5 (2), there exists a completely null-homologous reflexive link L_0 in M such that a handlebody can be obtained from M by a $1/\mathbb{Z}$-Dehn surgery along L_0. Moreover, by the proof of Lemma 2.5 (2), we can take L_0 so that $L_0 \cap \bigcup F_i = \emptyset$. Thus the completely disjoint spanning surface system $\{F_i\}$ is contained in the resultant handlebody V_0 obtained from M by a $1/\mathbb{Z}$-Dehn surgery along L_0. □

References

1. R. Benedetti, R. Frigerio, *Levels of knotting of spatial handlebodies*, Trans. Amer. Math. Soc. **365** (2013), 2099–2167.
2. F. Bonahon, *Cobordism of automorphisms of surfaces*, Ann. Sci. Éc. Norm. Supér. (4) **16** (1983), 237–270.
3. R. H. Fox, *On the imbedding of polyhedra in 3-space*, Ann. of Math. **49** (1948), 462–470.
4. J. Hempel, *A surface in S^3 is tame if it can be deformed into each complementary domain*, Trans. Amer. Math. Soc. **111** (1964), 273–287.
5. J. Hempel, 3*-manifolds*, Ann. of Math. Stud. **86**, Princeton, New Jersey, Princeton University Press, 1976.
6. W. Jaco, *Nonretractable cubes-with-holes*, Michigan Math. J. **18** (1971), 193–201.
7. W. Jaco. and D. R. McMillan, *Retracting three-manifolds onto finite graphs*, Illinois J. Math. **14** (1970), 150–158.
8. L. H. Kauffman, *On knots*, Princeton Univ. Press, Princeton 1987.
9. H. W. Lambert, *Mapping cubes with holes onto cubes with handles*, Illinois J. Math. **13** (1969), 606–615.

10. F. Lei, *Complete systems of surfaces in 3-manifolds*, Proc. Cambridge Phil. Soc. **122** (1997), 185–191.
11. H. Matsuda, M. Ozawa and K. Shimokawa, *On non-simple reflexive links*, J. Knot Theory and its Ramifications **11**, 787 (2002).
12. J. Morgan and G. Tian, *Ricci flow and Poincaré conjecture*, Clay Mathematics Monographs Vol.3, 2007.
13. K. Nakamura, *Fox reimbedding and Bing submanifolds*, to appear in Trans. Amer. Math. Soc..
14. G. Perelman, *The Entropy Formula for the Ricci Flow and Its Geometric Application* (2002), arXiv:math/0211159.
15. G. Perelman, *Ricci Flow with Surgery on Three-Manifolds* (2003), arXiv:math/0303109.
16. M. Scharlemann and A. Thompson, *Surfaces, submanifolds, and aligned Fox reimbedding in non-Haken 3-manifolds*, Proc. Amer. Math. Soc. **133** (2005), 1573–1580.
17. H. Seifert, *Über das Geschlecht von Knoten*, Math. Annalen **110** (1934), 571–592.

Department of Natural Sciences, Faculty of Arts and Sciences, Komazawa University, 1-23-1 Komazawa, Setagaya-ku, Tokyo, 154-8525, Japan
E-mail address: w3c@komazawa-u.ac.jp

Department of Mathematics, Saitama University, Saitama 338-8570, Japan
E-mail address: kshimoka@rimath.saitama-u.ac.jp